



IBD BENCHMARKING TOOL 2019/2020 RESULTS OVERVIEW

KENT AND CANTERBURY, WILLIAM HARVEY AND
QUEEN ELIZABETH THE QUEEN MOTHER HOSPITALS
(PAEDIATRIC SERVICE)

www.ibduk.org

This report shows results from the IBD Benchmarking Tool 2019/2020, a new way to assess how well services across the UK are providing care against the IBD Standards 2019. It includes information from a self-assessment completed by the IBD Team. Unfortunately, no information can be included from the IBD Patient Survey due to insufficient numbers of patients responding who use this service.

This report has been produced by IBD UK, a partnership of patient and professional organisations working together for everyone affected by IBD.

Key



IBD Patient Survey 2019



IBD Service Self-Assessment 2019/2020



The Service in Numbers

Covers a population of

720,500

Supports

50

paediatric IBD
patients


Of which

9

were diagnosed in the
past 12 months



The IBD Team



Quality of Care

Overall, how would you rate the quality of your Crohn's or Colitis care over the last 12 months?

Unfortunately, no information can be included here due to insufficient numbers of patients (less than 5) responding to the IBD Patient Survey for this service.

These results have been made publicly available to support clinicians, patients and commissioners to improve quality of care for IBD patients. Thank you to everyone who responded to the IBD Patient Survey 2019. The next IBD Patient Survey will run in 2021.



Patient Journey Overview

Key



IBD Patient Survey 2019



IBD Service Self-Assessment 2019/2020



Pre-diagnosis

All* patients with IBD are
seen within 4 weeks from
their first referral

[Achieved by 75% of services]



Not yet achieved

Newly Diagnosed



% of newly
diagnosed patients are
offered written information
on their condition

[Insufficient data: less than 5 responses]
[All UK survey responses: 96%]



% of patients were given
clear information about next steps
and who to contact while waiting
for their diagnosis to be confirmed

[Insufficient data: less than 5 responses]
[All UK survey responses: 64%]

New diagnoses and the
care plan that has been
agreed are communicated
to GPs within 1 week

[Achieved by 59% of services]



Achieved



Flare Management

All* IBD patients contacting the
IBD advice line get a response
by the end of the next
working day

[Achieved by 86% of services]



Not yet achieved



Guidance about steroid use is
communicated to primary care
and steroid use recorded

Not yet achieved

[Achieved by 59% of services]



All* IBD patients have access to
review by the IBD team within a
maximum of 5 working days

Achieved

[Achieved by 94% of services]

Data is routinely collected
and audited on patients
receiving more than
1 course of steroids a year

[Achieved by 12% of services]



Not yet achieved

All* inpatients with IBD are seen
by a consultant gastroenterologist
and/or colorectal surgeon
on a specialist ward within
24 hours of admission

[Achieved by 86% of services]



Not yet achieved



All* elective IBD
surgery takes place
within 18 weeks

Achieved

[Achieved by 91% of services]

Surgery and Inpatient Care





Patient Journey Overview

Key



IBD Patient Survey 2019



IBD Service Self-Assessment 2019/2020



Ongoing Care and Monitoring



% of patients have
contact with an IBD nurse
specialist

[Insufficient data: less than 5 responses]
[All UK survey responses: 97%]

% of patients have
regular reviews regardless
of their health

[Insufficient data: less than 5 responses]
[All UK survey responses: 84%]



All* IBD patients have a
personalised care plan based on
a holistic needs assessment
[Achieved by 60% of services]



Not yet achieved



% of patients believe
their care is well-coordinated
between their GP and
gastroenterologist

[Insufficient data: less than 5 responses]
[All UK survey responses: 47%]

% of patients discuss
wider life and priorities, as
part of planning their care

[Insufficient data: less than 5 responses]
[All UK survey responses: 47%]



The IBD Service

All* patients have access to
non-acute endoscopy and
imaging within 4 weeks, and
within 24 hours for patients who
are acutely unwell or require
admission to hospital



Achieved

[Achieved by 63% of services]

IBD patients are directly
involved in service
development



[Achieved by 20% of services]

Not yet achieved



% of patients have
been given the opportunity
to feedback on their care in
the past 12 months

[Insufficient data: less than 5 responses]
[All UK survey responses: 30%]



Protocols are in place to
support transition to adult
services for young people
with IBD

Not yet achieved

[Achieved by 78% of services]

All* patients with confirmed
IBD are recorded in an
electronic clinical
management system



[Achieved by 74% of services]

Not yet achieved